



Bowel Preparation Instructions – SUPREP

Any questions please call (631)-289-0300 ext. 224.

3 DAYS PRIOR to the PROCEDURE:

- Do not eat foods high in fiber.
- Examples are whole grain breads and cereals, fruits, nuts, seeds, quinoa, popcorn, and cooked or raw vegetables.
- Do not take any fiber supplements.

On the DAY BEFORE Your PROCEDURE:

DIET

- Consume only clear liquids the entire day before
- Okay to consume one protein shake, before 11 am, but only from the following lists:
 - Protein powder mixed with water or milk
 - Pre-made Protein shake
 - AVOID FRUIT or VEGETABLE shakes.

At 11 AM: Begin Drinking ONLY clear liquids. NO MORE PROTEIN SHAKES.

A clear liquid is one that can you see through. Examples of clear liquid are as follows:

- Clear beef or chicken broth
- Popsicles, except for red, purple or blue
- Coconut water (NOT coconut milk)
- Strained fruit juice such as apple or white grape juice
- Coffee and tea (without cream or dairy products)
- · Gatorade
- Soft drinks
- · Kool-Aid
- · Jell-O
- Do not drink alcohol

7pm - Start 1st Dose:

- 1. Pour ONE (1) 6 oz bottle of SUPREP[®] liquid into the mixing container.
- 2. Add cool drinking water to the 16 oz line on the container and stir to mix.
- 3. Drink ALL the liquid in the container.
- 4. You must drink two (2) more 16 oz containers of water over the next 1 hour.

9pm - start 2nd dose

- Follow steps 1 4 above.
- · Starting at midnight, you may have nothing by mouth, including water
- Drink plenty of clear liquids during and after bowel-prep process to help prevent dehydration

On the Morning/Day of your PROCEDURE

PREP – Procedure before 9am

- If you take daily medication(s), you may do so up until 2 hours before your scheduled report time with small sips of water.
- If you take blood thinners, follow the instructions your health care provider gives you.
- Arrange for a ride home. You will be drowsy after the exam and will need an escort, 18 years or older. If taking a medical cab service, you will need an escort as well to take you home.

The importance of a GOOD PREP

Your prep is one of the most important parts of your colonoscopy. Without a good prep, your bowel empty and clean, your gastroenterologist will not be able to see your colon clearly. This could lead to missing a polyp/lesions or rescheduling your colonoscopy.

PREP – Procedure 9am or later

7pm - Start 1st Dose:

- 1. Pour ONE (1) 6 oz bottle of $SUPREP^{\mathbb{R}}$ liquid into the mixing container.
- 2. Add cool drinking water to the 16 oz line on the container and stir to mix.
- **3**. Drink **ALL** the liquid in the container.
- 4. You must drink two (2) more 16 oz containers of water over the next 1 hour.
- Drink plenty of clear liquids during and after bowel-prep process to help prevent dehydration

On the Morning/Day of your PROCEDURE

6am - start 2nd Dose:

- Follow steps 1 4 above.
- Be sure to finish drinking all of your laxative by 7am.
- If you take daily medication(s), you may do so up until 2 hours before your scheduled report time with small sips of water.
- If you take blood thinners, follow the instructions your health care provider gives you.
- Arrange for a ride home. You will be drowsy after the exam and will need an escort, 18 years or older. If taking a medical cab service, you will need an escort as well to take you home.

The importance of a GOOD PREP

Your prep is one of the most important parts of your colonoscopy. Without a good prep, your bowel empty and clean, your gastroenterologist will not be able to see your colon clearly. This could lead to missing a polyp/lesions or rescheduling your colonoscopy.